



ACC.14

TCT@ACC-12 | innovation in intervention

A202

JACC April 1, 2014

Volume 63, Issue 12



## Acute Coronary Syndromes

### CAN A SECOND MEASUREMENT OF COPEPTIN IMPROVE ACUTE MYOCARDIAL INFARCTION RULE OUT?

Poster Contributions

Hall C

Sunday, March 30, 2014, 3:45 p.m.-4:30 p.m.

Session Title: Acute Coronary Syndromes: Treatment Considerations

Abstract Category: 1. Acute Coronary Syndromes: Clinical

Presentation Number: 1226-242

Authors: Nicholas Marston, Kevin Shah, Christian Mueller, Sean-Xavier Neath, Robert Christenson, James McCord, Richard Nowak, Gary Vilke, Lori Daniels, Judd Hollander, Fred Apple, Chad Cannon, John Nagurney, Donald Schreiber, Christopher DeFilippi, Christopher Hogan, Deborah Diercks, Alexander Limkakeng, Inderjit Anand, Allan Jaffe, W. Frank Peacock, Alan Maisel, Alan Wu, University of California San Diego, San Diego, CA, USA

**Background:** Management of chest pain patients with an initial negative troponin but an elevated copeptin level is unclear. We hypothesized that a second copeptin measurement two hours after presentation would help risk-stratify these patients.

**Methods:** The multi-center CHOPIN trial enrolled 2071 acute chest pain patients. 494 subjects with an initial negative troponin and elevated copeptin ( $>14$  pmol/L) were included in this study. Copeptin and troponin levels were rechecked at 2 hours and the final diagnosis of acute myocardial infarction (AMI) was made by two independent, blinded cardiologists.

**Results:** Of the 494 patients analyzed, 378 (76.5%) had a persistently elevated copeptin at 2 hours and 116 patients (23.5%) had a copeptin that fell below the cutoff of 14 pmol/L. The AMI rate was 6.1% when the second copeptin was elevated compared to 0% when the second copeptin was not ( $p=.006$ ), yielding a sensitivity of 100%. This strategy identified 23.5% of patients with an intermediate risk who could be ruled out for AMI while still in the ED. In contrast, a second troponin measurement had a sensitivity of 74%.

**Conclusions:** A negative second copeptin drawn 2 hours after presentation demonstrated 100% sensitivity for AMI, allowing for effective rule out in this intermediate risk cohort. This strategy identified nearly 25% of intermediate risk patients who could be considered for discharge, creating the potential for a significant reduction in the number of unnecessary chest pain admissions.

**Figure: Second Copeptin vs. Second Troponin for AMI Rule Out in initial Trop-/Cop+**

	2nd Copeptin	2 <sup>nd</sup> Troponin
Sensitivity for AMI	100% (95% CI 85.0 to 100%)	73.9% (95% CI 51.6 to 89.7%)
Negative Predictive Value	100% (95% CI: 96.8 to 100%)	98.7% (95% CI: 97.16 to 99.5%)
Negative Likelihood Ratio	.00 (95% CI: not applicable)	.28 (95% CI: .14 to .55)